



Wildflower Information & Planting Instructions

Mixture Formulation : Annuals have been included to establish cover quickly and to give color the first year; some may produce new plants the following year (the biennials may also reseed). Perennial plants live for more than two years, and most flower from the second year onward. Very few wildflowers bloom continually throughout the season; therefore, we have included spring-, summer-, and fall-blooming species in each mixture. Colors include blue, purple, red, white, yellow and pink. Mix heights vary from 10 inches to 8 feet.

Site Selection : Does the site support plants now? If you have a site where nothing, including weeds, is growing, that site is unlikely to support wildflowers. Will there be adequate moisture during germination and establishment? Can you supply supplemental water, if necessary? What weed seeds are likely to be present in the soil? Will weeds spread to your site from adjacent areas?

Planting Rates : Each mixture has a recommended minimum and maximum planting rate. The planting rates for some mixtures are adjusted higher due to the presence of small-seeded, nonaggressive species. In general however, the minimum planting rates are based on 60-70 seeds per square foot (4 to 12 pounds per acre), which is usually sufficient to establish a good stand of wildflowers on prepared soil when adequate weed control can be maintained. Maximum planting rates are based on 120-140 seeds per square foot (8 to 24 pounds per acre), and are recommended when adequate soil preparation and weed control are not possible, or when maximum color is required. Avoid using more than the recommended rates since poor perennial establishment may result.

When to Plant : The best time to plant in your area depends on the climate and rainfall patterns as well as the species you are planting. In cool climates, plant annuals, perennials or mixtures of annuals and perennials in spring, early summer or late fall. Fall plantings should be late enough so that seeds do not germinate until spring. Perennials can also be sown in early fall provided that there are at least 10-12 weeks of growing time before the plants go dormant for the winter. Late fall plantings are advantageous when supplemental irrigation cannot be provided and adequate rainfall is anticipated in the spring.

Use of Grasses : Wildflowers can be sown alone or with grasses. For most areas of the United States, we recommend using native grass mixtures. Aggressive grasses should be avoided because they will crowd out most wildflowers; these grasses include Kentucky Bluegrass, Smooth Brome, Crested Wheatgrass, Bermuda Grass, and Annual Rye. If wildflowers must be used with these grasses, the flowers should be planted in high-density patches as accents to the grassed areas. Or the flowers may be sown with the grasses if the planting rates of the grasses are reduced significantly.

Site Preparation : Proper site preparation is important for prompt germination of seed and healthy growth of seedlings. Best results will be obtained by planting on cleared ground. Remove existing vegetation to avoid competition from other plants. This may be done by pulling, tilling under, spraying with a general herbicide, or by a combination of these methods, depending upon the size of the area, type and density of vegetation and other factors. Loosen soil by scraping, tilling or scarifying. Tilling should be utilized only when soil is very compacted and further weed control measures can be taken. Specific recommendations are given under the heading, Weed Control.

Seed Application : Method of application depends on the size of the area and the terrain. On small areas, broadcast seeds evenly either by hand or by use of a drop or cyclone spreader. It is helpful to mix a carrier such as clean, dry sand with the seed; sand adds volume and aids in even distribution. We recommend using a ratio of 1 or 2 parts sand to 1 part seed. Rake in lightly, covering seeds to a maximum depth of 2-3 times their thickness. Or drag the area lightly with a piece of chain link fence to mix the seed into the surface of the soil.

Moisture : Best results will be obtained by soaking the planted areas thoroughly and maintaining consistent moisture for 4-6 weeks - then gradually reducing waterings. In non-irrigated situations, plant in the spring or before periods of anticipated rainfall. After seedlings are established, watering may be reduced depending on the climate and rainfall. In arid climates or during drought conditions, up to 1/2 inch of supplemental water per week may be required to maintain an optimal display. If weeds are present, remember that they benefit from moisture as much as the wildflowers and may dominate overwatered areas.

Fertilization : Many wildflowers benefit from some fertilization if the soil does not have adequate nutrients. Some wildflowers do fine in poor soils, while others require a more fertile environment. We recommend that a soil test be performed when soil quality is unknown. If the soil needs improvement, use a low nitrogen fertilizer with a 5-10-10 ratio or add organic matter such as weed-free straw or grass clippings, well-rotted compost, peat moss, or leaf mold. In addition to adding nutrients, organic materials enhance soil structure and encourage beneficial microorganisms. Avoid over-fertilizing which may promote weed growth and lush foliage rather than flowers.

Weed Control : Weed control is the biggest problem facing plant establishment and one which has no easy solution. Weed seeds are present in many situations and lie dormant, but viable, for long periods. A weedy area converted to wildflowers will have a large reservoir of weed seeds in the soil, ready to germinate when conditions are favorable. In most cases, it is advisable to consider weed control in two phases-as part of site preparation prior to planting, and as an important component of the post-germination maintenance program. Before planting, remove existing weeds by pulling, tilling under, applying a glyphosate herbicide such as Roundup®, or by a combination of these methods. For additional weed control after site preparation, a soil fumigant may be used, or the area may be irrigated to encourage weed growth and then sprayed with a general herbicide. In very weedy areas, the following method is suggested: (1) Till soil or spray vegetation with Roundup®. When using an herbicide, allow vegetation to die, then rake out the dead debris. If perennial weeds such as bindweed are present, using an herbicide is more effective than tilling. (2) Irrigate to encourage germination of weed seeds near the surface; most seeds will germinate within two weeks if consistent moisture is available. Do not till the soil again because this will bring even more weed seeds up to the surface. (3) Spray any new growth with Roundup®. (4) After raking out dead vegetation, allow soil to recover for 3-4 weeks before planting seed. From our experience, a recovery period of this duration is advisable because extensive use of glyphosate herbicides may cause a delay in germination and in the vigorous growth of seedlings. Once the seeds have germinated, further weed control is usually necessary. If practical, pull all weeds as soon as they can be identified. Other successful techniques are spot-spraying with a general herbicide or selectively cutting weeds with a string trimmer. Be sure to remove weeds before they reseed. Many unwanted annual and some perennial grasses can be controlled with the herbicides Grass-B-Gon*, Ornamec* and Fusilade*. These post-emergents do not affect broad-leaved plants so they can be applied over existing flowers; they are most effective when sprayed on new growth and young plants. Take care to avoid treating areas with desirable native grasses or fescues. *Observe all precautions and follow manufacturer's recommendations for application.

What to Expect : Avoid seeding more than the recommended rate since overseeding can result in crowded conditions the first year and poor establishment of perennials. Cover seeds lightly to protect them from drying out during germination, and to prevent them from being eaten by birds. Consistent moisture is important for 4 - 6 weeks after planting. A wildflower planting requires the same weed control measures as traditional landscaping. Effective measures include site preparation prior to planting and a post-germination maintenance program. Wildflower mixes contain annual, biennial and perennial species. The annuals, which may not be native to your area, are included to assure maximum color during the first season and to act as a nurse crop for the slower-growing perennials. Annuals germinate quickly when conditions are favorable, providing a quick ground cover and competition against weeds. Natural reseeding of annuals ranges from significant to minimal, depending on the species, climate, soil texture and other factors. Most perennial and biennial species begin to bloom the second season, but not as profusely as annuals. Therefore, wildflower plantings look noticeably different after the first year. Sometimes it is desirable or even necessary to sow seed in second and subsequent years. Reseeding may be necessary if establishment of wildflowers is spotty or poor. It is possible to reseed bare areas with the original mixture. Loosen soil of bare areas and provide adequate weed control and supplemental irrigation as needed. Where natural reseeding of annuals is minimal, sowing annuals each spring can produce a magnificent annual and perennial display throughout the growing season. If desired, wildflowers may be mowed in the fall following seed set. Mow to a height of 4-6 inches, and leave the residue on the ground because it is a reservoir of viable seeds.

Bird & Butterfly Wildflower Mix: Seeding rate: 10 - 20 pounds per acre or 1 1/4-2 1/2 pounds per 5,000 square feet.
15% Lupine, Arroyo – 15% Mallow, Tree – 15% Four-O'clock – 15% Nasturtium – 8% Zinnia 'Pumila Mix' – 5% Birds Eye – 4% Larkspur, Rocket – 4% Pinks, Sweet William – 3% Snapdragon, Spurred – 3% Poppy, Corn – 3% Sage, Scarlet – 3% Catchfly – 2% Balsam – 1% Foxglove – 1% Maltese Cross – 1% Mint, Lemon – 1% Penstemon, Rocky Mountain

Deer Proof Wildflower Mix: Seeding rate: 10 - 20 pounds per acre or 1 1/4 - 2 1/2 pounds per 5,000 square feet.
22.24% Lupine, Arroyo – 17.79% Lupine, dwarf 'Pixie Delight' - 13.34% Lupine, Perennial – 8.89% Poppy, California – 8.89% Gaillardia, Perennial – 6.67% Sage, Scarlet – 6.67% Zinnia 'Pumila Mix' – 4.45% Lupine, Yellow – 4.45% Poppy, corn – 2.22% Coreopsis, Plains - 2.22% Alyssum, Tall White Sweet – 1.11% Daisy, Gloriosa – 0.56% Sage, Blue – 0.27% Yarrow, White – 0.22% Foxglove

Partial Shade Wildflower Mix: Seeding rate: 5 - 10 pounds per acre or 3/4 - 1 1/4 pounds per 5,000 square feet.
22.00% Forget-Me-Not, Chinese - 15.00% Candytuft, Annual - 12.00% Cornflower, Blue - 11.00% Baby Blue-Eyes - 10.00% Five-Spot - 7.00% Coreopsis, Lance Leaved - 5.00% Pinks, Sweet William - 4.00% Daisy, Shasta - 4.00% Larkspur, Rocket - 3.00% Clarkia - 2.00% Snapdragon, Spurred - 2.00% Poppy, Corn - 1.00% Columbine, Mixed Colors Giant - 1.00% Foxglove - 1.00% Alyssum, Tall White Sweet

Perennial Wildflower Mix: Seeding rate: 5 - 10 pounds per acre or 3/4 - 1 1/4 pounds per 5,000 square feet.
22.0% Wallflower, Siberian - 17.0% Coreopsis, Lance-leaved - 16.0% Flax, Blue - 11.0% Lupine, Perennial - 11.0% Foxglove - 5.5% Gaillardia, Perennial - 5.0% Pinks, Sweet William - 5.0% Daisy, Shasta - 2.0% Columbine, Dwarf - 1.5% Hyssop, Lavender - 1.0% Evening, Primrose, Dwarf - 0.50% Daisy, Gloriosa - 0.50% Aster, New England - 0.50% Coneflower, Prairie - 0.50% Mexican Hat - 0.50% Black-eyed Susan - 0.50% Daisy, English